
Agenda

- Reminder: Observation Thursday night
- · Pass back graded stuff
- Class discussion: Scientific Argument
- Ch. 7

My Theory

- Moon is a hollow spherical world constructed by an advanced alien civilization
- · Defend or attack



The Solar System


- The *Solar System* is occupied by a diversity of objects, but shows an underlying order in the dynamics of their movements
- The planets form two main families:
 - solid rocky inner planets
 - gaseous/liquid outer planets
- From these observations, astronomers deduce that the Solar System formed some 4.5 billion years ago out of the collapse of a huge cloud of gas and dust




- The Sun is a star, a ball of incandescent gas whose output is generated by nuclear reactions in its core
- Composed mainly of hydrogen (71%) and helium (27%), it also contains traces of nearly all the other chemical elements

The Sun


• It is the most massive object in the Solar System – 700 times the mass of the rest of the Solar System combined





 Its large mass provides the gravitational force to hold all the Solar System bodies in their orbital patterns around the Sun




The Planets • All of the planets travel counterclockwise around the Sun (as seen from high above the Earth's north pole) • Six planets rotate counterclockwise; Venus rotates clockwise (retrograde rotation), and Uranus and Pluto appear to rotate on their sides









Dwarf Planets

- Pluto and similar objects fail to fit into either family
- Recently, scientists have discovered more than 200 similar objects orbiting the Sun at the same distance as Pluto
- In 2006, a new family was introduced the dwarf planets
 - Massive enough to pull themselves spherical
 - Orbits have not been swept clear of debris

Satellites

- The number of planetary satellites changes frequently as more are discovered!
 - Jupiter 62
 - Saturn 31
 - Uranus 27
 - Neptune 13
 - Mars 2
 - Earth 1
 - Mercury and Venus are moonless







Asteroids and Comets


Typical comet orbi

- Composition and size
 - Asteroids are rocky or metallic bodies ranging in size from a few meters to 1000 km across (about 1/10 the Earth's diameter)
 - Comets are icy bodies about 10 km or less across that can grow very long tails of gas and dust as they near the Sun and are vaporized by its heat

Asteroids and Comets

- Their location within Solar System
 - Most asteroids are in the asteroid belt between
 Mars and Jupiter indicating that these asteroids are the failed building-blocks of a planet
 - Some comets may also come from a disk-like swarm of icy objects that lies beyond Neptune and extends to perhaps 1000 AU, a region called the Kuiper Belt

Asteroids and Comets



Most comets orbit the Sun far beyond Pluto in the *Oort cloud*, a spherical shell extending from 40,000 to 100,000 AU from the Sun

Measuring Composition

- Since the inner and outer planets differ dramatically in composition, it is important to understand how composition is determined
- A planet's reflection spectrum can reveal a planet's atmospheric contents and the nature of surface rocks
- Seismic activity has only been measured on Earth for the purposes of determining interior composition

Measuring Composition: Density




- A planet's average density is determined by dividing a planet's mass by its volume
 - Mass determined from Kepler's modified third law
 - Volume derived from a planet's measured radius

Measuring Composition: Density


- Once average density known, the following factors are taken into account to determine a planet's interior composition and structure:
 - Densities of abundant, candidate materials
 - Variation of these densities as a result of compression due to gravity
 - Surface composition determined from reflection spectra
 - Material separation by density differentiation
 - Mathematical analysis of equatorial bulges

Analysis Concludes:



 The terrestrial planets, with average densities ranging from 3.9 to 5.5 g/cm³, are largely rock and iron, have iron cores, and have relative element ratios similar to the Sun except for deficiencies in lightweight gasses

Analysis Concludes:



 The Jovian planets, with average densities ranging from 0.71 to 1.67 g/cm³, have relative element ratios similar to the Sun and have Earth-sized rocky cores

Age of the Solar System

- All objects in the Solar System seem to have formed at nearly the same time, out of the same original cloud of gas and dust
- Radioactive dating of rocks from the Earth, Moon, and some asteroids suggests an age of about 4.5 billion yrs
- A similar age is found for the Sun based on current observations and nuclear reaction rates

Bode's Law

- First noted in 1766, formalized mathematically by J. E. Bode in 1778
 - 0 3 6 12 24 48 96 192 384
 - 4 7 10 16 28 52 100 196 388
 - 0.4 0.7 1.0 1.6 2.8 5.2 10.0 19.6 38.8
- Does a pretty good job, up to a point



Origin of the Solar System

- A theory of the Solar System's formation must account for the following:
 - Planets orbit in the same direction and in the same plane
 - Rocky inner planets and gaseous/liquid/icy outer planets
 - Compositional trends in the solar system
 - All Solar System bodies appear to be less than 4.5 billion years old
 - Other details structure of asteroids, cratering of planetary surfaces, detailed chemical composition of surface rocks and atmospheres, etc.

The Solar Nebula Hypothesis

- Derived from 18th century ideas of Laplace and Kant
- Proposes that Solar System evolved from a rotating, flattened disk of gas and dust (an interstellar cloud), the outer part of the disk becoming the planets and the inner part becoming the Sun



The Solar Nebula Hypothesis

- This hypothesis naturally explains the Solar System's flatness and the common direction of motion of the planets around the Sun
- Interstellar clouds are common between the stars in our galaxy and this suggests that most stars may have planets around them



Interstellar Clouds

- Come in many shapes and sizes – one that formed Solar System was probably a few light years in diameter and 2 solar masses
- Typical clouds are 71% hydrogen, 27% helium, and traces of the other elements




Interstellar Clouds

- Clouds also contain tiny dust particles called *interstellar* grains
 - Grain size from large molecules to a few micrometers
 - They are a mixture of silicates, iron and carbon compounds, and water ice




In the Beginning...



 Triggered by a collision with another cloud or a nearby exploding star, rotation forces clouds to gravitationally collapse into a rotating disk

The Solar Nebula



- A few million years pass for a cloud to collapse into a rotating disk with a bulge in the center
- This disk, about 200 AU across and 10 AU thick, is called the solar nebula with the bulge becoming the Sun and the disk condensing into planets

Temperatures in the Solar Nebula

- Before the planets formed, the inner part of the disk was hot, heated by gas falling onto the disk and a young Sun – the outer disk was colder than the freezing point of water
- · Gas/dust disks have been observed

Ice-rich planetesimals

Disk Observations





Condensation

Condensation
 occurs when gas
 cools below a
 critical temperature
 at a given gas
 pressure and its
 molecules bind
 together to form
 liquid/solid particles




Condensation in the Solar Nebula

- Iron vapor will condense at 1300 K, silicates will condense at 1200 K, and water vapor will condense at room temperature in air
- In a mixture of gases, materials with the highest vaporization temperature condense first
- Condensation ceases when the temperature never drops low enough
- Sun kept inner solar nebula (out to almost Jupiter's orbit) too hot for anything but iron and silicate materials to condense
- Outer solar nebula cold enough for ice to condense


Formation of Planets Copyright O The Midrae-Hill Comparins. Inc. Permission required for reproduction of display. Solar rediction heets up the linear solar system. Water-ice only in couler parts of disk Iron and silicate dust particles throughout disk Iron and silicate-rich dust planetesimals Outer parts of disk Kuiper belt Inner planets

Accretion

- Next step is for the tiny particles to stick together, perhaps by electrical forces, into bigger pieces in a process called accretion
- As long as collisions are not too violent, accretion leads to objects, called planetesimals, ranging in size from millimeters to kilometers




Planetesimals



- Planetesimals in the inner solar nebula were rockyiron composites, while planetesimals in the outer solar nebula were icyrocky-iron composites
- Planets formed from "gentle" collisions of the planetesimals, which dominated over more violent shattering collisions


Formation of the Planets

- Simulations show that planetesimal collisions gradually lead to approximately circular planetary orbits
- As planetesimals grew in size and mass their increased gravitational attraction helped them grow faster into clumps and rings surrounding the Sun



Formation of the Planets

- Planet growth was especially fast in the outer solar nebula due to:
 - Larger volume of material to draw upon
 - Larger objects (bigger than Earth) could start gravitationally capturing gases like H and He



Continuous Bombardment



Lanks



• Continued planetesimal bombardment and internal radioactivity melted the planets and led to the density differentiation of planetary interiors


Formation of Moons







- Moons of the outer planets were probably formed from planetesimals orbiting the growing planets
- Not large enough to capture H or He, the outer moons are mainly rock and ice giving them solid surfaces




Formation of Atmospheres

- Atmospheres were the last planet-forming process
- Outer planets gravitationally captured their atmospheres from the solar nebula
- Inner planets created their atmospheres by volcanic activity and perhaps from comets and asteroids that vaporized on impact
- Objects like Mercury and the Moon are too small

 not enough gravity to retain any gases on their surfaces

Extra-solar Planets

- Evidence exists for planets around other nearby stars
- The new planets are not observed directly, but rather by their gravitational effects on their parent star
- These new planets are a surprise - they have huge planets very close to their parent stars



Extra-solar Planets

- Idea: The huge planets formed far from their stars as current theory would project, but their orbits subsequently shrank
- This migration of planets may be caused by interactions between forming planets and leftover gas and dust in the disk

